arXiv:1004.2973v1 [math.AP] 17 Apr 2010

In this note we prove that:

Theorem 1. *for* $2 \leq s < \frac{n}{2}$ *or* $1 \leq s < \frac{2n}{n+1}$ *or* $1 \leq s < \frac{n}{2}$ *but n is even,*

$(-\Delta)^s(u) = |u|^{q-2}u, q = \frac{2n}{n-2s}$

has infinitely many sign changing solutions or equivalently we can say that there exist solutions u_k *such that* $\int u_k(-\Delta)^s(u_k)dx \to \infty$ *as* $k \to \infty$

A brief history of this problem: In 1979, Gidas,Ni and Nirenberg[4] classified all the positive solutions when $s = 1$. In 1999, Wei and Xu [5]classified the positive solutions when s is integer. In 2004 and 2006, Li[9] and Chen,Li and Ou [6]classified the positive solutions for $0 < s < \frac{n}{2}$. In 1986, Ding[1] proved the above theorem for case $s = 1$,In 2004 Bartsch, Schneider and Weth [3]proved the cases when s is integer. The ideas used are same , both pull back the energy functional to sphere and then use symmetric criticality to obtain the critical points.

Proof of the theorem:

In the following , dx always denotes the volume form of R^n , $d\xi$ denotes the volume form of sphere S^n with standard metric , c denotes some constant.

First notice that the solutions of the equation in the theorem are exactly the critical points of the functional $J(u) = \frac{1}{2}\int u(-\Delta)^s u dx - \frac{1}{q}\int |u|^q dx$

Step 1:

Use stereographic projection to lift $(-\Delta)^s$ to sphere

$\pi : R^n \longrightarrow S^n$, namely

$\pi(x) = (\frac{2x}{1+|x|^2}, \frac{1-|x|^2}{1+|x|^2})$

By direct computation we have that the Jacobian is given:

$J_\pi = (\frac{2}{1+|x|^2})^n = cU^{\frac{2n}{n-2s}} = cU^q$.

Then the intertwining operator A_s on S^n ([8])is defined:

$A_s(w) \circ \pi = cJ_\pi^{-\frac{n+2s}{2n}}(-\Delta)^s(J_\pi^{\frac{n-2s}{2n}}(w \circ \pi)) = cU^{1-q}(-\Delta)^s(U(w \circ \pi))$

The eigenvalues of A_s are given by $\lambda_l = \frac{\Gamma(\frac{n}{2}+l+s)}{\Gamma(\frac{n}{2}+l-s)}$, the corresponding eigenspaces are spanned by orthonormal spherical harmonics (same with the standard laplacian on S^n).

Consider the map:

$\Theta : H^s(R^n) \to H^s(S^n)$,

$\Theta(v) = (U^{-1}v) \circ \pi^{-1}$

where $H^s(R^n)$ is the completion of compact supported functions under norm $\sqrt{((-\Delta)^s(\cdot), \cdot)}$ and $H^s(S^n)$ is the completion of smooth functions under norm $\sqrt{(A_s(\cdot), \cdot)}$

Direct computations show that Θ is an isometry and it also preserves the L_q norm so we have:

$\tilde{J}(\Theta(v)) := \frac{1}{2}\int \Theta(v)A_s(\Theta(v))d\xi - \frac{1}{q}\int |\Theta(v)|^q d\xi = J(v) = \frac{1}{2}\int v(-\Delta)^s v dx - \frac{1}{q}\int |v|^q dx$

Which means that the critical points of $\tilde{J}$ and J are in 1-1 correspondence.

Step2:

Now we study the critical points of $\tilde{J}$.

Since $\tilde{J}$ is invariant under conformal transform of S^n , which means that it does not satisfy the Palais-Smale condition.However, applying as in [1] or [3] the symmetric mountain pass theorem [7] and the principle of symmetric criticality [2] we have the following.

Lemma1. *Let G be a compact subgroup of $O(n+1)$ acting linearly and isometrically on $H^s(S^n)$ such that*

(1) $\tilde{J}$ is G-invariant;

(2) the embedding $H^s_G(S^n) \hookrightarrow L^q(S^n)$ is compact;

(3) $H^s_G(S^n)$ has infinite dimension.

Then $\tilde{J}$ has a sequence of critical points w_k , such that $\int w_k A_s(w_k) d\xi \to \infty$ as $k \to \infty$.

Here we denote by $H^s_G(S^n)$ the subspace of $H^s(S^n)$ consisting of G-invariant functions: $H^s_G(S^n) = \{w \in H^s(S^n) | w(g\xi) = w(\xi)$, every $g \in G$ and a.e. $\xi \in S^n\}$

(1) and (3) in Lemma1 are obvious, in the next step we will check (2) for some G.

Step3:

Now let $G = O(\lfloor \frac{n}{2} \rfloor) \times O(\lfloor \frac{n+1}{2} \rfloor)$ (here ($\lfloor \frac{n}{2} \rfloor$) means the greatest integer less or equal than $\frac{n}{2}$),let $m < s < m+1$,m is an integer.It is easy to see that the minimum dimension of the orbit of G is $d_G := \lfloor \frac{n}{2} \rfloor$.

Elementary computations show that $q = \frac{2n}{n-2s} < \frac{2(n-d_G)}{n-d_G-2m}$, apply Lemma3.2 in [3] we see that the embedding $H_G^m(S^n) \hookrightarrow L^q(S^n)$ is compact.

Then from the formula of eigenvalues of A_s it is easy to see that $\sqrt{(A_s(\cdot),\cdot)}$ is increasing as s increases, which means that we have a continuous embedding $H_G^s(S^n) \hookrightarrow H_G^m(S^n)$

So the embedding $H_G^s(S^n) \hookrightarrow L^q(S^n)$ is compact and it satisfies (3) in Lemma1 .

Finally , by applying Lemma1 we prove Theorem1.

Reference:

[1] Ding, W., On a conformally invariant elliptic equation, Comm. Math. Phys. 107 (1986), 331-335.

[2] Palais, R. S., The principle of symmetric criticality, Comm. Math. Phys. 69 (1979), 19-30.

[3] T. Bartsch, M. Schneider, T. Weth; Multiple solutions to a critical polyharmonic equation, J. Reine Angew. Math. 571(2004), 131-143 .

[4] Gidas, B., Ni, W.-M., and Nirenberg, L., Symmetry and related properties via the maximum principle, Comm. Math. Phys. 68 (1979), 209-243.

[5] Wei, J., and Xu, X., Classification of solutions of higher order conformally invariant equations, Math. Ann. 313 (1999), 207-228.

[6] W. Chen, C. Li, and B. Ou, Classification of solutions for an integral equation, Comm. Pure Appl. Math., 59 (2006), pp. 330-343.

[7] Ambrosetti, A., Rabinowitz, P.H.: Dual variational methods in critical point theory and applications. J. Funct. Anal. 14, 349-381 (1973)

[8] C. Morpurgo, Sharp trace inequalities for intertwining operators on Sn and Rn. Internat. Math. Res. Notices (1999), 20, 1101-1117.

[9]Li, Y.Y.: Remark on some conformally invariant integral equations: the method of moving spheres. J. Eur. Math. Soc. 6, 153-180 (2004)